

3-D SIMULATIONS OF THE CHEMICAL AND DYNAMICAL EVOLUTION OF THE GALACTIC BULGE

NAOHITO NAKASATO^{1,2} AND KEN'ICHI NOMOTO¹
Appear in the Astrophysical Journal

ABSTRACT

A three-dimensional hydrodynamical N-body model for the formation of the Galaxy is presented with special attention to the formation of the bulge component. Since all previous numerical models for the Galaxy formation do not have a proper treatment of the chemical evolution and/or sufficient spatial resolutions, we have constructed a detailed model of the chemical and dynamical evolution of the Galaxy using our GRAPE-SPH code. Our SPH code includes various physical processes related to the formation of stellar systems. Starting with cosmologically motivated initial conditions, we obtain a qualitatively similar stellar system to the Galaxy. Then we analyze the chemical and kinematic properties of the bulge stars in our model and find qualitative agreement with observational data. The early evolution of our model has revealed that most bulge stars form during the sub-galactic merger (merger component of the bulge stars). Because of the strong star burst induced by the merger, the metallicity distribution function of such stars becomes as wide as observed. We find that another group of the bulge stars forms later in the inner region of the disk (non-merger component of the bulge stars). Because of the difference in the formation epoch, the main source of iron for this group of stars is different from the merger component. Iron in the merger and non-merger components comes mainly from Type II and Type Ia supernovae, respectively. Since a Type Ia supernova ejects ~ 10 times more iron than a Type II supernova, $[\text{Fe}/\text{H}]$ of the non-merger component tends to be higher than that of the merger component, which widens the metallicity distribution function. From these results, we suggest that the Galactic bulge consists of two chemically different components; one has formed quickly through the sub-galactic clump merger in the proto-galaxy and the other has formed gradually in the inner disk.

Subject headings: Milky Way: Bulge — galaxy: formation — stars: formation — hydrodynamics

1. INTRODUCTION

The formation of bulges, which are the high density central component of spiral galaxies, is a key process in the formation of spiral galaxies. There are several scenarios for the formation process of the bulges (e.g., Bouwens, Cayon, Silk 1999). Such scenarios can be divided into the following categories; (1) a primordial collapse (e.g., Eggen, Lynden-Bell, & Sandage 1962), (2) merging of sub-clumps in a proto-galactic cloud (e.g., Baugh, Cole, & Frenk 1996), and (3) a secular evolution of a stellar disk (e.g., Norman, Sellwood, & Hasan 1996). Previously, the surface brightness of the bulges has been recognized to have the $R^{1/4}$ profile as in ellipticals, where R denotes the distance from the center (e.g., Whitford 1978; Rich 1988). However, recent observations have revealed the presence of a class of bulges that are well fitted by the exponential profile (e.g., Andredakis, Peletier, & Balcells 1995), i.e., there exist two classes of bulges according to the surface brightness. Each formation scenario has advantage and disadvantage to explain the diversity of bulges (Wyse 1999).

For the formation of the Galactic bulge, Matteucci & Brocato (1990) and Matteucci, Romano, & Molaro (1999) have constructed one-zone chemical evolution models as follows: (1) They assumed that the Galactic bulge was formed by a single star-burst. In their best fit model, the formation time scale is ~ 0.1 Gyr. (2) One of the predictions of their calculations is that the element ratio such as

$[\text{O}/\text{Fe}]$ remains high even for metal-rich stars ($[\text{Fe}/\text{H}] > 0$) because the formation time scale of the bulge is shorter than the lifetime of a Type Ia supernova progenitor. However, the chemical properties predicted by the one-zone model depend strongly on their *assumption* of the dynamics, i.e., the formation time-scale of the bulge. To constrain the formation scenarios, we need to calculate the chemical and dynamical evolution together and compare the model predictions with both the chemical and kinematic properties of the bulge.

Here, we present our first results of the three-dimensional Galactic bulge formation model. We model the formation of the Galaxy by means of Smoothed Particle Hydrodynamics (SPH) method (Lucy 1977; Gingold & Monaghan 1977) that incorporates a star formation algorithm and a self-consistent treatment of the chemical enrichment history of gas. The three-dimensional models for the formation of disk galaxies have been investigated by many authors (e.g., Katz 1992; Steinmetz & Müller 1995; Friedli & Benz 1995; Berczik 1999) and they have succeeded in many respects. We follow the method and models by the previous authors but use much larger number of particles to study the detailed formation and evolution processes of the Galaxy (Nakasato 2000).

In Cold Dark Matter (CDM) cosmology, the early stage of the formation of galaxies involves the progressive merging of sub-galactic clumps (e.g., Baugh, Cole, & Frenk 1996). To properly model the dynamics of the merger

¹ Department of Astronomy, School of Science, University of Tokyo, Bunkyo-ku, Tokyo 113-0033; nakasato@astron.s.u-tokyo.ac.jp, nomoto@astron.s.u-tokyo.ac.jp

² Astronomisches Rechen-Institut, Heidelberg, 69120

history, we adopt the three-dimensional SPH method. The chemodynamical approach by Samland, Hensler, & Theis (1997), which is a mesh based and two-dimensional method, is not suitable to model the early merging history of the proto-galaxy.

Our SPH code uses the GRAPE (Sugimoto et al. 1990) and the Remote-GRAPE system (Nakasato, Mori, & Nomoto 1997) to compute the gravitational interaction between particles (GRAPE-SPH code). The adoption of the Remote-GRAPE system doubles the performance of our code since we can make the SPH calculation and N-body summation in parallel (Nakasato, Mori, & Nomoto 1997). This enhancement of performance enables us to use seven times larger number of particles than Katz (1992), Steinmetz & Müller (1995), and Raiteri, Villata, & Navarro (1996) so that the spatial resolution of our model is almost two times higher than their models. As a result, we can investigate the early merging history in much more detail during the Galaxy formation. We note that other authors (e.g., Navarro & Steinmetz 2000; Sommer-Larsen & Dolgov 2001) have published the comparable or even higher resolution SPH galaxy formation models than the present work; however, these authors have mainly concentrated on the kinematic properties of the models. In contrast, in the present work, we have made attention to both the chemical and kinematic properties.

Our study is the first attempt to model the formation and chemical evolution of the Galactic bulge using the self-consistent three-dimensional SPH method. Using the chemical and dynamical SPH code, we simulate the evolution of the Galaxy starting from a plausible cosmological initial condition. The obtained model well reproduces the important chemical and kinematic properties of the Galaxy such as the formation of the bulge, disk and halo as seen in Figure 5. By analyzing the result of this model, we construct a plausible scenario for the Galactic bulge formation.

The plan for the present paper is as follows: In section 2, we briefly describe our SPH method. Section 3 describes the procedure to construct initial conditions. We present a summary of results obtained with our model in section 4. By analyzing the result of our Galaxy model, we describe the formation history of the bulge component in section 5, and compare the chemical and kinematic properties of the bulge with recent observational results. Finally, section 6 is devoted to conclusions.

2. THE METHOD

We adopt the SPH method to construct self-consistent three-dimensional dynamical and chemical models for the formation of stellar systems. The SPH method has been applied to many astrophysical problems. Because of its Lagrangian nature, it is suitable to systems that have large density contrasts, e.g., the formation of galaxies (e.g., Evrard 1988; Hernquist & Katz 1989; Katz 1992; Steinmetz & Müller 1994), the evolution of galaxies (e.g., Friedli & Benz 1995; Patsis & Athanassoula 2000), the cosmological simulations (e.g., Navarro & White 1994; Yoshikawa, Jing, & Suto 2000), and a cloud-cloud collision (e.g., Lattanzio et al. 1985; Habe & Ohta 1992). Various codes have been developed to combine SPH and N-body systems. In these codes, gravitational forces are calculated

with various methods such as direct summations, Particle-Particle/Particle-Mesh methods (e.g., Evrard 1988), Tree methods (e.g., Hernquist & Katz 1989; Benz et al. 1990), and the method to use the special purpose computer GRAPE (e.g., Umemura et al. 1993; Steinmetz 1996).

To simulate the formation of a stellar system from gas, we use our GRAPE-SPH code using the Remote-GRAPE library (Nakasato, Mori, & Nomoto 1997). The SPH formulation that we use is the same as Navarro & White (1993). We use a spatially variable smoothing length and integrate equations of motion with a second order Runge-Kutta method as described in Navarro & White (1993). To simulate the formation of stellar systems from gas, our GRAPE-SPH code includes various physical processes related to the formation of stellar systems, e.g., cooling, star formation, and feedback by stars as follows.

2.1. Cooling and star formation

We adopt the metallicity dependent cooling functions, which are computed by MAPPINGS III package by R.S. Sutherland (MAPPINGS III is the successor of MAPPINGS II described in Sutherland & Dopita 1993), to solve the energy equation of gas particles (see Figure 1 of Nakasato, Mori & Nomoto (2000) for the actually used cooling functions)

For the star formation algorithm, we adopt the star formation recipe as used in the usual SPH codes (Katz 1992). Hereafter, “STAR” means “star particle”, which is not an individual star but an association of many stars. In this recipe, we convert a fraction of a gas particle into a STAR when four conditions are satisfied. The first three conditions are the *physical* conditions expressed as

$$(\nabla \cdot \mathbf{v})_i < 0, \quad (1)$$

$$t_{\text{cool}} < t_{\text{d}}, \quad (2)$$

$$t_{\text{d}} < t_{\text{sound}}, \quad (3)$$

where t_{cool} , t_{d} , and t_{sound} are the cooling time, dynamical time, and sound crossing time of the gas particle, respectively (see Nakasato, Mori & Nomoto (2000) for the detailed time scale definition).

During each star formation interval ($\delta t = 2$ Myr in the present work), we check whether the three conditions are satisfied for each gas particle. If the gas particle satisfies these three conditions, we then check the forth *probability* condition. Assuming star formation process is a Poisson process (Katz 1992), we can write the probability for star formation within δt as

$$P = 1.0 - \exp(-\delta t/t_{\text{starform}}), \quad (4)$$

where t_{starform} is the local star formation time scale. In our SPH code, we define $t_{\text{starform}} = t_{\text{d}}/C$, where C is a parameter to control our star formation recipe. If P is larger than a random number ($0 - 1$), we create a new STAR from a fraction of the gas particle (star formation occurs!). Introducing this *probability* condition, we can effectively control the threshold density for star formation by changing the parameter C . We compute P as a function of the density (ρ) for different values of C as shown in Figure 1. Here, we assume $\delta t = 2$ Myr. Note that the timestep used in the time integration is always smaller than δt . From Figure 1, we can see effects of C as follows; larger (smaller) C means lower (higher) threshold density.

To demonstrate the dependence on the star formation parameter C , we evolve a spherical region of $10^{12} M_{\odot}$, which consists of 10 % gas and 90 % dark matter, for various C . Initially, we set up the sphere in rigid rotation and outward expansion. And the angular speed of rotation corresponds to the spin parameter $\lambda \sim 0.05$, where $\lambda = \frac{L|E|^{1/2}}{GM^{5/2}}$, with L being the angular momentum, E the total energy, M the mass, and G the gravitational constant (Padmanabhan 1993). This initial condition is the same as used in Katz (1992) and Steinmetz & Müller (1995). We evolve the sphere for three different star formation parameters $C = 0.1, 0.5$ and 1.0 . In Figure 2, we present the star formation history for $C = 0.1$ and $C = 1.0$. For $C = 1.0$, the star formation rates (SFR) increase more rapidly than $C = 0.1$ since the threshold density for star formation is lower. These STARS which formed early (i.e., very old STARS) in the $C = 1.0$ model have large vertical velocities. If we compare the projected edge-on surface density of STARS, the $C = 1.0$ model clearly shows more extended distribution than the $C = 0.1$ model. In Table 2, some kinematic properties at $t = 5$ Gyr are compared for different C . There is a clear correlation between C and the ratio between the X and Z axis velocity-dispersion (or the spin parameter). From these results, we conclude that the $C = 0.1$ and $C = 1.0$ galaxies are similar to late type and early type galaxies, respectively. To summarize, if we use this *probability* condition, various values of C lead to various global star formation histories. Hereafter, we set $C = 0.1$ throughout the present paper.

The mass of a newly formed STAR (M_{star}) is calculated as

$$M_{\text{star}} = \left[1 - \frac{1}{1 + 0.5 \frac{\delta t}{t_{\text{starform}}}} \right] \pi h^3 \rho, \quad (5)$$

where h and ρ are the smoothing length and the density of the gas particle, respectively (Nakasato, Mori & Nomoto 2000). In the present calculation, the typical mass of STARS is $\sim 10^6 - 10^7 M_{\odot}$.

2.2. Feedback from stars

Formed stars eject energy and mass as stellar winds and supernova explosions (feedback from stars). As a result, stars heat up, accelerate, and enrich the circumstellar and interstellar medium. With current computing resources, it is not yet feasible to accurately include the energy, momentum, and mass feedback from stars in our SPH code. This is because the length resolution of the present calculation (~ 0.5 kpc) is much longer than a typical size of supernova remnants (< 100 pc). Thus, in the present paper, we adopt a phenomenological method as adopted by several authors (e.g., Katz 1992) to mimic real feedback processes; i.e., we distribute the energy and mass ejected by a STAR to the neighbor gas particles. In our SPH code, we incorporate the effect of stellar winds and Type II and Type Ia supernova explosions as the energy and mass feedback processes. We compute the mass and energy ejection rates based on a simple model explained below. In this paper, we distribute the feedback energy as purely thermal energy. This treatment is a simple and a zero-th order approximation to the real nature and other authors have tried to refine the treatment of feedback (e.g., Thacker & Couchman 2000; Springel & Hernquist 2002).

Also in this paper, we follow the chemical evolution of total metal (Z), iron (Fe) and oxygen (O). The chemical evolution is coupled with the thermal and dynamical evolution, i.e., the ejected metal affects the evolution of gas through the metallicity dependent cooling rates. To summarize, our GRAPE-SPH code is a most up-to-date SPH implementation for the formation of stellar systems and used to successfully model the formation of globular clusters (Nakasato, Mori & Nomoto 2000).

2.3. Energy ejection

The energy ejection rate per STAR is given as

$$E_{\text{eject}} = e_{\text{SW}} R_{\text{SW}} + e_{\text{SNII}} R_{\text{SNII}} + e_{\text{SNIa}} R_{\text{SNIa}}, \quad (6)$$

where e_{SW} is the total ejected energy by stellar winds during the stellar life time and e_{SNII} and e_{SNIa} are the energies ejected by a Type II and Ia supernova explosion, respectively. The rate R_{SW} is the number of stars that expel their envelopes at the current epoch per unit time and R_{SNII} and R_{SNIa} are the rates of Type II and Type Ia supernovae, respectively. We define R_{SW} and R_{SNII} as follows

$$R_{\text{SW}} = \frac{\int_{M_{\text{ms}}}^{M_{\text{up}}} \phi(m) dm}{\tau(M_{\text{ms}})} \quad (7)$$

$$R_{\text{SNII}} = \frac{\int_{M_{\text{ms}}}^{M_{\text{ma}}} \phi(m) dm}{\tau(M_{\text{ms}}) - \tau(M_{\text{ma}})}, \quad (8)$$

where $\phi(m)$ is the initial mass function (IMF), namely $\phi(m)dm$ gives the number of stars in the mass range of $(m, m + dm)$, and $\tau(m)$ is the stellar lifetime as a function of stellar mass (David, Forman & Jones 1990). In the present study, we assume the power-law type IMF as

$$\phi(m) = A m^{-2.35}, \quad (9)$$

where A is a constant. For the upper and lower mass limits in the IMF, $M_{\text{up}} = 120 M_{\odot}$ and $M_{\text{lo}} = 0.05 M_{\odot}$ are assumed. In Eqs. (7) and (8), $M_{\text{ma}} (= 50 M_{\odot})$ and $M_{\text{ms}} (= 8 M_{\odot})$ are the upper and lower mass limits of the stars that explode as Type II supernovae.

For the Type Ia supernova rate, we adopt the formulation given in the galactic chemical evolution model by Kobayashi et al. (1998, 2000). They adopt the single-degenerate scenario (Nomoto et al. 1994) for the Type Ia supernova progenitor, which is that a white dwarf (WD) in a close binary undergoes a thermonuclear explosion when the companion star evolves off the main-sequence and transfers a large enough amount of mass over to the WD. In their model, the Type Ia supernova rate at an epoch t is given as

$$R_{\text{SNIa}}(t) = C_{\text{SNIa}} \frac{\int_{M(t)}^{M(t+\Delta t)} \phi(m) dm}{\Delta t}, \quad (10)$$

where $M(t)$ is the mass of the companion star whose main-sequence lifetime is t and C_{SNIa} is a constant to be calibrated by the observational constraints. From the evolution model of the progenitor stars, Hachisu, Kato & Nomoto (1996, 1999) obtained the mass ranges of the

companion stars of the WDs that become Type Ia supernovae as

$$M_{\text{sec}} = [1.8, 2.6], \quad [0.9, 1.5] \quad M_{\odot}. \quad (11)$$

We note that these mass ranges weakly depend on the metallicity of the star. In the present paper, we don't adopt the metallicity dependence. Figure 3 shows R_{SNIa} as a function of time, where we set $C_{\text{SNIa}} = 1.0$. In the present paper, we set $C_{\text{SNIa}} = 4.0 \times 10^{-4}$ from the result of test calculations. Note that other authors (Raiteri, Villata, & Navarro 1996; Carraro, Lia & Chiosi 1998) adopt different progenitor models by Greggio & Renzini (1983) and Matteucci & Greggio (1986).

For the supernova energy, we assume that $e_{\text{SNI}} = e_{\text{SNIa}} = 10^{51}$ erg. For the stellar wind energy, e_{SW} is estimated to be 0.2×10^{51} erg for solar metallicity stars from the observational data of OB associations (Abbot 1982). The chemical abundance of a massive star significantly affects e_{SW} (Leitherer, Robert, & Drissen 1992), so that we use metallicity dependent e_{SW} as $e_{\text{SW}} = 0.2e_{\text{SNI}}(Z/Z_{\odot})^{0.8}$, where Z is the mass fraction of heavy elements in the STAR.

2.4. Mass ejection

In our code, the mass ejection in stellar winds from massive stars ($m \geq M_{\text{ms}}$) is combined with the mass ejection due to Type II supernovae. The mass ejection in stellar winds from low mass stars ($m < M_{\text{ms}}$) is treated separately. Thus, the mass ejection rate per STAR is written as

$$M_{\text{eject}} = m_{\text{SNI}}R_{\text{SNI}} + m_{\text{SWm}}R_{\text{SWm}} + m_{\text{SNIa}}R_{\text{SNIa}}, \quad (12)$$

where m_{SNI} is the average ejecta mass of Type II supernovae, and m_{SWm} is the average mass ejected in stellar winds from the low mass stars (corresponding to planetary nebulae), and m_{SNIa} is the ejecta mass of a Type Ia supernova. R_{SWm} is the number of stars per unit time expelling their envelopes at a certain epoch. m_{SNI} and m_{SWm} are defined as

$$m_{\text{SNI}} = \frac{\int_{M_{\text{ms}}}^{M_{\text{ma}}} m\phi(m)dm}{\int_{M_{\text{ms}}}^{M_{\text{ma}}} \phi(m)dm} - m_{\text{NS}}, \quad (13)$$

$$m_{\text{SWm}} = \frac{\int_{M_{\text{II}}}^{M_{\text{ms}}} m\phi(m)dm}{\int_{M_{\text{II}}}^{M_{\text{ms}}} \phi(m)dm} - m_{\text{WD}}. \quad (14)$$

Here m_{NS} is the mass locked up in a neutron star and m_{WD} is the mass that is locked up in a WD. $M_{\text{II}} (= 1 M_{\odot})$ is the mass of the star whose life time nearly equals to the Hubble time. We set $m_{\text{NS}} = 1.4 M_{\odot}$ and $m_{\text{WD}} = 1.4 M_{\odot}$ and assume that a Type Ia supernova is produced by a Chandrasekhar mass WD so that we set $m_{\text{SNIa}} = 1.4 M_{\odot}$. We define R_{SWm} as

$$R_{\text{SWm}} = \frac{\int_{M_{\text{II}}}^{M_{\text{ms}}} \phi(m)dm}{\tau(M_{\text{II}}) - \tau(M_{\text{ms}})}. \quad (15)$$

The fraction of heavy elements in m_{SNI} and m_{SNIa} is computed using the nucleosynthesis yield of Type II and Ia

supernovae (Tsujimoto et al. 1996; Nomoto et al. 1997). We compute the chemical evolution of total metal (Z), iron (Fe) and oxygen (O) in our code. The metallicity yield we used is tabulated in Table 1.

2.5. Summary on Feedback

The feedback phase is divided into three phases; a stellar wind phase, a Type II supernova phase, and a Type Ia supernova phase.

- The stellar wind phase continues for $\tau(M_{\text{ma}})$, during which only the energy ejection from STARS is included; the ejected mass is included in the Type II supernova phase for simplicity.
- The Type II supernova phase begins at $t = \tau(M_{\text{ma}})$ and ends at $t = \tau(M_{\text{ms}})$. During this phase, the mass ejection rate is the sum of the contributions by stellar winds of massive stars and Type II supernovae.
- The Type Ia supernova phase begins at $t = \tau(M_{\text{ms}})$. During this phase, both the energy ejection and mass ejection rates are the sum of the contributions by Type Ia supernovae and stellar winds of low mass stars.

We present the schematic view of the feedback processes in Figure 4.

With the present algorithm and the adopted parameters, a STAR of $10^8 M_{\odot}$ produces $\sim 5.5 \times 10^5$ Type II supernova explosions and $\sim 1.4 \times 10^5$ Type Ia supernova explosions during 13 Gyr of the evolution. Also $\sim 2.2 \times 10^7 M_{\odot}$ gas is ejected during the evolution. The ejected gas contains $\sim 1.6 \times 10^6 M_{\odot}$ of heavy elements, which include $\sim 1.5 \times 10^5 M_{\odot}$ of iron and $\sim 1.0 \times 10^6 M_{\odot}$ of oxygen.

As a final note, the thermal energy, gas, and heavy elements from stellar winds and supernovae are smoothed over neighbor particles of the STAR within a neighbor radius of R_f (a feedback radius). R_f is a parameter set to 0.5 kpc, which is equal to the adopted gravitational softening length of STAR particles. Results of a test calculation with $R_f = 0.8$ kpc are very similar to those of the present paper.

3. INITIAL MODEL

Following the previous work (e.g., Katz 1992), we investigate the evolution of the spherical 3σ top hat over-dense region of a mixture of dark matter and gas with our SPH code. We neglect the matter outside the spherical regions, thus neglecting the late in-fall of the external matter and the external gravitational field. Throughout the paper, we set the Hubble constant $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega = 1$, and $\Lambda = 0$. We set the initial redshift (z) of the model to be ~ 25 and the initial radius of the sphere to be 55 kpc. Thus, the comoving radius of the spherical region is $\sim 1.4 \text{ Mpc}$ and the contained mass is $\sim 10^{12} M_{\odot}$. We set the region in rigid rotation to provide a sufficient angular momentum with the spin parameter $\lambda \sim 0.1$. We also add the corresponding Hubble velocity to the velocity field of the sphere since we integrate the model with the physical coordinate.

To generate a 3σ over-dense region, we use the path integral method (Bertschinger 1987). We note that there are further two methods to set up the over-dense region: (1) To seek the desired over-dense region by sampling different random field realizations, which is used by the cosmological hydrodynamical simulations of formation of X-ray clusters (Anninos & Norman 1996). (2) To select the desired halo from the results of the large scale cosmological N-body simulations and use it as an initial condition for a cosmological galaxy formation model (Navarro & White 1994). The path integral method is the fastest method among other methods. And it is the easiest way to generate a several dozen desired over-dense regions, i.e. the over-dense region of $10^{12} M_\odot$ in the present study. The weakness of this method is that we have to use the isolated boundary condition. Namely, we neglect the external tidal field and artificially add the angular momentum to the initial velocity field. On the other hand, with the method (2), it is natural to use the surrounding particles as the origin of external tidal-field. However, with the method (2), it is difficult to find several over-dense regions with similar properties since each region is very different like the real universe. We think our way to generate initial conditions is an efficient way for our purpose to construct the model for our Galaxy.

Specifically, initial models are constructed as follows: (1) We generate a spherical top hat 3σ over-dense region by the path integral method (Bertschinger 1987). We use the COSMICS package (Bertschinger 1995) and generate 50 realizations. (2) We follow the non-linear evolution of the dark matter particles in each spherical region with the same code. Note that we only use the N-body part of our GRAPE-SPH code in these simulations. (3) At a certain epoch ($z \sim 2.5$), we examine the properties of dark matter halos and select an appropriate realization for the hydrodynamical model. We select the model with a single halo since if two dominant halos exist, the gas disk will be destroyed by a major merger event (Barnes & Hernquist 1992). Among 50 realizations, 39 realizations are single halo models and others are multiple halo models. From 39 realizations, we select candidates for the hydrodynamical model by comparing the model density profile with the density profile of the Galactic halo. (4) Once we select several models, we restart the hydrodynamical N-body simulation from $z \sim 2.5$ with our GRAPE-SPH code. After we have done full hydrodynamical calculations for three single-halo models (and two multiple-halo models), we select the best-fit model, which is presented in this paper, by comparing the disk scale length with the Galactic scale length. All results shown in the following sections are obtained with this best-fit model.

We have computed a similar kind of models with several different configurations and found that the performance mainly depends on the speed of the GRAPE system: (a) For our old configuration using four GRAPE-3A boards, a similar calculation to the present work takes more than 200 hours up to $z \sim 0.9$ ($t = 5$ Gyr). (b) For our recent configuration using one GRAPE-5 board, a similar calculation takes only ~ 150 hours up to $z = 0$ ($t = 13$ Gyr). Using this recent configuration, a full hydrodynamical N-body computation for several dozen realizations are under way and results will be presented elsewhere.

4. FORMATION OF THE GALAXY

The initial number of gas particles and dark matter particles are $\sim 27,000$ and $\sim 27,000$, respectively. Thus, with the total mass of $10^{12} M_\odot$, the initial masses of the gas particles and dark matter particles are $4 \times 10^6 M_\odot$ and $3.6 \times 10^7 M_\odot$, respectively. We set the gravitational softening length for the gas and dark matter particles to be 0.5 kpc and 1.0 kpc, respectively. The overall evolution is very similar to the previously reported work (Steinmetz & Müller 1995; Berczik 1999). During the expansion due to the Hubble flow, small scale structures grow and dense regions collapse. In such high density regions (clumps), star formation occurs as a result of the efficient radiative cooling. These small clumps gradually merge to produce larger clumps. At $t \sim 0.3 - 0.5$ Gyr, these clumps merge and the resulted clump becomes the primary halo that eventually becomes a spiral galaxy. Around this time ($z \sim 7$), the star formation rates (SFR) reaches a maximum value. At $t \sim 1$ Gyr ($z \sim 4$), the dark matter is almost virialized and the evolution of dark matter particles becomes a quasi-static state while the gas particles are being settled into a thin disk. Finally, the prominent gas disk forms in the center of the primary halo at $t \sim 3$ Gyr ($z \sim 1.6$). Afterward, the most star formation occurs in this gas disk and the overall evolution becomes a quasi-static state.

After 5 Gyr of evolution (up to $z \sim 0.9$), over 80,000 STARS have formed and the number of gas particles have decreased to $\sim 13,000$. At this stage, we categorize STARS according to the chemical properties and the formation epoch (t_{form}) of STARS following the observational characteristics of the three components of the Galaxy as defined in Mihalas & Binney (1981).

Specifically, we categorize STARS by the following conditions and show the projected stellar densities in Figure 5:

- left panel : Metal-poor ($\log(Z/Z_\odot) < -3$) STARS,
- center panel : Old ($t_{\text{form}} < 1$ Gyr) and metal-rich ($\log(Z/Z_\odot) > 0$) STARS,
- right panel : Young ($t_{\text{form}} > 4$ Gyr) STARS.

Here, Z is the mass fraction of heavy elements in STARS. Metal-poor STARS show an extended distribution since these STARS formed at an early epoch and at high latitude (the left panel). The old and metal-rich STARS are concentrated in the galactic center (the center panel). The qualitative properties of the system are not so changed after $z \sim 1$ as reported by Steinmetz & Müller (1995). Since we neglect the late in-fall of surrounding material, the most recent star formation mainly occurs in the disk. As a result, the young STARS are located near the galactic plane (the right panel). These results show that the calculated stellar system looks very similar to the Galaxy. To be more qualitatively, we plot the face-on surface density profile of STARS in Figure 7. The solid crosses show our result. We fit the model with the following function; $\rho(r) = \rho_{\text{disk}} \exp(-r/R_{\text{disk}}) + \rho_{\text{bulge}} \exp(-7.67((r/R_{\text{bulge}})^{0.25} - 1))$, where the scale radii are $R_{\text{disk}} = 4.21$ kpc and $R_{\text{bulge}} = 0.876$ kpc. In the next section, we analyze the properties of STARS near the galactic center.

At $z = 0$, the total number of particles has increased to $\sim 150,000$ where the numbers of gas, dark matter, and

STAR particles are $\sim 11,000$, $27,000$, and $112,000$, respectively. When we compare the projected density profile at $z \sim 0.9$ (Figure 5) and at $z = 0$, the 3 components structure looks very similar except the thickness of the disk STARS. Most star formation occurs quasi-statically near the disk plane and the SFR is $\sim 1 - 2 M_{\odot} \text{ yr}^{-1}$.

5. FORMATION HISTORY OF THE BULGE COMPONENT

The one-zone chemical evolution model (e.g., Matteucci, Romano, & Molaro 1999) predicted the approximate chemical properties of the Galactic bulge. To construct a self-consistent model of the bulge formation, however, a chemical and dynamical model is required. In this section, we use our high resolution model for the formation and evolution of the Galaxy to study the formation of the Galactic bulge.

5.1. Dynamical history and kinematics of the bulge stars

We select the STARS located near the galactic center ($R < 2 \text{ kpc}$) and define these STARS as the bulge stars. We define the center of the gravity of all STARS as the galactic center. To find the galactic center, we need several times of a iteration to compute the center of the gravity. In the i -th iteration, we compute the center of the gravity of STARS that is located within $R_i \text{ kpc}$ from the origin of the coordinate and convert the origin with that point. We set $R_1 = 20 \text{ kpc}$ and $R_2, R_3, \dots = 5 \text{ kpc}$. Typically, we only need three or four iterations. For example, at $z \sim 0.9$, the mass and number of the bulge stars are $1.8 \times 10^{10} M_{\odot}$ and $16,800$, respectively. Ever after this time, we can select the bulge stars with the same definition. In this paper, however, we compare our model results at this time with some observational data. The reason is that in SPH models including star formation recipes, the resolution of the model becomes worse and worse during the evolution since the number of gas particles decreases. With our chemical enrichment model, we need to have a large enough number of gas particles around a STAR particle to properly model the chemical evolution of the interstellar matter; further, the gas particles show a more extended distribution than the star particles and this distribution makes the effective resolution of the chemical enrichment description worse. At $z \sim 0.9$, the number of gas particles becomes $\sim 13,300$, which is a half of the initial number of gas particles and sufficiently large.

Figure 6 shows the SFR as a function of time for the bulge stars (solid line) and all stars (dotted line) up to $z \sim 0.9$. In this figure, we normalize the SFR by the total mass of each component at $z \sim 0.9$. More than 60 % of the bulge stars up to this time formed during the first 0.5 Gyr of the evolution. With our model, we can obtain the detailed formation history of the bulge stars. In Figure 8, we present snap shots for projected STAR positions in the region where star formation is most violent for the first 0.5 Gyr. We note that the merging of two sub-galactic clumps occurs during $t = 0.4 - 0.5 \text{ Gyr}$. This merging of the clumps causes the star burst of the short time scale shown in Figure 6. Such a phenomenon is different from the simple star burst assumed in the one-zone chemical evolution model (Matteucci & Brocato 1990).

With our chemical and dynamical model, we can also obtain the kinematics of the bulge stars. In Figure 9, we plot

the average velocity (upper line) and the velocity dispersion (lower line) of these bulge stars. The bulge stars are substantially rotationally supported with the rotational speeds of $\sim 160 - 230 \text{ km s}^{-1}$. These speeds are smaller than the rotational speeds of the disk stars ($\sim 270 \text{ km s}^{-1}$ in the present model), which means that the angular momentum of the bulge is lost during the merging process. Figure 10 shows the x-direction velocity dispersion for the bulge stars as a function of $[\text{Fe}/\text{H}]$. Here, our numerical results are shown by the solid triangles and solid squares and compared with observational data (Minniti 1996) shown by the dotted circles (note that the observational data is the line of sight velocity dispersion). The solid squares are the STARS with $t_{\text{form}} < 1 \text{ Gyr}$ (old STARS) and the solid triangles are the STARS with $t_{\text{form}} < 5 \text{ Gyr}$ (young STARS). The old and young STARS show a different trend, and the result for old STARS is more consistent with the observational data. Our model (especially old STARS) qualitatively reproduces the kinematic properties of the Galactic bulge.

5.2. Chemical properties of bulge stars

Important information to constrain the formation of the Galactic bulge concerns the chemical properties of the bulge stars. In this subsection, we analyze the chemical properties of the bulge stars.

The observed metallicity distribution function of K-giant stars (McWilliam & Rich 1994) is characterized by a wide ($-1.2 < [\text{Fe}/\text{H}] < 0.9$) distribution as shown by the dashed line in Figure 11. The distribution function of the bulge stars in our model (the solid line in Figure 11) is as wide as $-2.0 < [\text{Fe}/\text{H}] < 1.0$, being in good agreement with the observational data. As noted in the previous subsection, about 60 % of the bulge stars form in the strong star-burst caused by merging of sub-clumps (hereafter these stars are called a merger component). This merger-driven star-burst is partly responsible for this wide distribution. On the other hand, other 40 % of the bulge stars form in the late stage of evolution (a non-merger component) and also contribute to widen the distribution function. This is because the non-merger component of the bulge stars has much larger $[\text{Fe}/\text{H}]$ than the merger component as explained below.

The bulge stars can be divided into two chemically different types with and without being contaminated by Type Ia supernova ejecta. With the adopted progenitor model (Kobayashi et al. 1998), the epoch of the first appearance of Type Ia supernovae $t_{\text{SNIa}} = 0.5 \text{ Gyr}$ after the formation of each STAR. Thus, the STARS that formed after $t = 0.5 \text{ Gyr}$ are partially contaminated by Type Ia supernovae. Each Type Ia supernova ejects ~ 10 times more iron than a Type II supernova (Nomoto et al. 1997). As a result, iron contained in the non-merger component of the bulge stars partially originates from Type Ia supernovae and their $[\text{Fe}/\text{H}]$ tends to be higher than that of the merger component. In Figure 12, the distribution function of the STARS that formed before and after $t = 0.5 \text{ Gyr}$ are plotted by the dashed and dotted lines, respectively, along with the total distribution function of the bulge stars (solid line). Both dashed and dotted lines show rather wide distribution and clearly the peak value of each distribution is different. The very old STARS, which form before $t =$

0.5 Gyr, contribute mainly to the metal poor range of the distribution.

The adopted yields are $[\text{O}/\text{Fe}] = 0.4$ from Type II supernovae and $[\text{O}/\text{Fe}] = -1.5$ from Type Ia supernovae (Table 1). Thus the contamination by Type Ia supernovae ejecta decreases $[\text{O}/\text{Fe}]$ in STARS starting from $[\text{O}/\text{Fe}] = 0.4$. In Figure 13, we present the metallicity distribution function of the bulge stars for $[\text{O}/\text{Fe}]$. The very peaky distribution shows that a half of the bulge stars have $[\text{O}/\text{Fe}] \sim 0.4$, i.e. a half of the bulge stars are not contaminated by Type Ia supernovae. Also, almost 80 % of the bulge stars have $[\text{O}/\text{Fe}] > 0$. Figure 14 shows $[\text{O}/\text{Fe}]$ as a function of $[\text{Fe}/\text{H}]$ for the bulge stars. $[\text{Fe}/\text{H}]$ increases up to ~ 0.3 without Type Ia supernova contamination. Later, due to the increasing Fe enrichment by Type Ia supernovae, $[\text{O}/\text{Fe}]$ decreases down to ~ -0.4 as $[\text{Fe}/\text{H}]$ increases to ~ 0.7 . We note for a given $[\text{Fe}/\text{H}]$ (> -0.5), $[\text{O}/\text{Fe}]$ is distributed over a wide range of > -0.4 . Thus, even for $[\text{Fe}/\text{H}] > 0$, large fraction of the bulge stars have $[\text{O}/\text{Fe}] > 0$, while $[\text{O}/\text{Fe}]$ is mostly sub-solar for $[\text{Fe}/\text{H}] > 0.3$. These features of our model are consistent with the observed results as plotted in Figure 14 by the dotted circles (Rich & McWilliam 2000). The observed $[\text{Si}/\text{Fe}]$ and $[\text{Ca}/\text{Fe}]$ show also relatively large dispersion from -0.4 to $+0.2$ for $[\text{Fe}/\text{H}] > 0$, although the observed results are still preliminary (Rich & McWilliam 2000).

In contrast, Matteucci & Brocato (1990) argued that the metallicity of the bulge stars mainly originated from Type II supernovae. In their model, they assumed $t_{\text{SNIa}} \sim 1$ Gyr and the star formation time scale is shorter than 1 Gyr so that the majority of Type Ia supernovae have exploded long after the star formation. Therefore the one-zone chemical evolution model predicts that the Galactic bulge is a chemically single population. In contrast to this picture, our chemical and dynamical model predicts that the Galactic bulge is neither a chemically single population and a dynamically single population.

5.3. Discussion

There are several problems in our numerical model. First, the present model, which is compared with some observational data, is just the current best-fit model from only 5 realizations. Thus, the quantitative comparisons between our model and observation are still far from complete. One reason is that it is too difficult to find a consistent initial model for very good match to the observation before doing real calculations. Due to complicated merging events occurring in the early phase, it is impossible to predict which initial halo will become a Milky-Way-like spiral galaxy.

Another point is the resolution of the numerical model. As seen in Figure 7, the inner region of the present model is not well resolved. Thus, our definition radius for the bulge, $R < 2$ kpc, is not a clear definition when we compare our results with observational data. This is because the Baade's window, where the chemical properties of the bulge stars are observed, is located 0.5 kpc from the galactic center. The choice of our definition is determined from the limitation of the numerical resolution of the present model. We use the rather large radius of 2 kpc to define the bulge stars since the gravitational softening length of the present model is 0.5 kpc (in fact, the inspection of Fig-

ure 7 shows that the effective resolution in our model is ~ 1 kpc, which equals to the softening length of the dark matter particles). To summaries, the number of particles in the present model is still too small to fully resolve the chemical structure of the bulge. To make more sensible comparison between numerical models and the observational data, we need much higher resolution models and more observational data obtained with large telescopes. Although there is such limitation, our model is in a good agreement with chemical and kinematic properties of the Galactic bulge stars.

6. CONCLUSION

In this paper, we report on the result of our high resolution three-dimensional models of the Galactic bulge formation. By modeling the formation and evolution of the Galaxy, we understand the formation history of the Galactic bulge as follows.

The early evolution of the best-fit model reveals that most bulge stars form during the merger that occurred in the region of the deepest potential. Because of the strong star burst induced by the merger, the metallicity distribution of the bulge component becomes as wide as observed. Although such a strong star burst was a similar effect as the burst assumed in the previous chemical evolution model, our model show that there is another reason to widen the metallicity distribution function of the bulge. By analyzing our results, we find that the bulge stars in our model are composed of two different components. One component (merger component) was produced by the merger occurred in the center of the halo as noted above. The other non-merger component was formed in the inner region of the disk after this merger. Due to the different formation epoch, the main source of iron for each component is different. Iron in the former comes mainly from Type II supernovae, and the other from Type Ia supernovae. Since each Type Ia supernova ejects ~ 10 times more iron than a Type II supernova, $[\text{Fe}/\text{H}]$ of the non-merger component tends to be higher than that of the merger component, which widened the metallicity distribution function. Also, our model predicts a relatively large dispersion in $[\text{O}/\text{Fe}]$ for $[\text{Fe}/\text{H}] > 0$.

Although our model is not in a complete agreement with our Galaxy, some chemical and kinematic properties of the Galactic bulge, such as the velocity dispersion vs. metallicity, are well reproduced. We thus conclude that an old fraction of the Galactic bulge is very likely to have been formed by the sub-clump merger in the proto-galactic environment, while a younger fraction of the Galactic bulge has formed gradually in the inner disk. These two groups should show different chemical properties.

We would like to thank the anonymous referee for the valuable comments and suggestions to greatly improve the manuscript. Also, we would like to thank J. Makino for kindly providing access to the GRAPE-4 and GRAPE-5 systems, on which a part of the computation has been done. Some part of the computation has been performed with the GRAPE cluster at the Astronomical Data Analysis Center of the National Astronomical Observatory, Japan. N.N. specially thanks the Astronomisches Rechen-Institut, Heidelberg for the hospitality during the stay as a guest researcher where a part of the paper has been

written. This work has been supported by JSPS Research Fellowships for Young Scientists (7664) and (05127), in part by an SFB 439 grant at the University of Heidelberg

and by the grant-in-Aid for Scientific Research (07CE2002, 12640233, 14047206, 14540223) of Ministry of Education, Science, Culture, Sports and Technology in Japan.

REFERENCES

- Abbot, D.C., 1982, *ApJ*, 263, 723
 Andredakis, Y.C., Peletier, R.F., & Balcells, M., 1995, *MNRAS*, 275, 874
 Anninos, P., & Norman, M., 1996, *ApJ*, 459, 12
 Benz, W., Bowers, R., Cameron, A., & Press, W., 1990, *ApJ*, 348, 647
 Berczik, P., 1999, *A&A*, 348, 371
 Barnes, J.E., & Hernquist L., 1992, *ARAA*, 705
 Baugh, C.M., Cole, S., & Frenk, C.S., 1996, *MNRAS*, 283, 1361
 Bertschinger, E., 1987, *ApJL*, 323, 103
 Bertschinger, E., 1995, <http://arcturus.mit.edu/cosmics/>
 Bouwens, R., Cayon, L., & Silk, J., 1999, *ApJ*, 516, 77
 Carraro, G., Lia, C., & Chosi, C., 1998, *MNRAS*, 297, 1021
 David, L.P., Forman, W., & Jones, C., 1990, *ApJ*, 459, 29
 Eggen, O.J., Lynden-Bell, D., & Sandage, A.R., 1962, *ApJ*, 136, 748
 Evvard, A. E., 1988, *MNRAS*, 235, 911
 Friedli, D., & Benz, W., 1995, *A&A*, 301, 649
 Gingold, A., & Monaghan, J., 1977, *MNRAS*, 181, 374
 Greggio, L., & Renzini, A., 1983, *A&A*, 118, 217
 Habe, A., & Ohta, K., 1992, *PASJ*, 44, 203
 Hachisu, I., Kato, M., & Nomoto, K., 1996, *ApJL*, 470, 97
 Hachisu, I., Kato, M., & Nomoto, K., 1999, *ApJ*, 522, 487
 Hernquist, L., & Katz, N., 1989, *ApJS*, 70, 419
 Katz, N., 1992, *ApJ*, 391, 502
 Kobayashi, C., Tsujimoto, T., Nomoto, K., Hachisu, I., & Kato, M., 1998, *ApJL*, 503, 155
 Kobayashi, C., Tsujimoto, T., & Nomoto, K., 2000, *ApJ*, 539, 26
 Lattanzio, C., Monaghan, J., Pongracic, H., & Schwarz, P., 1985, *MNRAS*, 215, 125
 Leitherer, C., Robert, C., & Drissen, L., 1992, *ApJ* 401, 596
 Lucy, L., 1977, *AJ*, 82, 1013
 Matteucci, F., & Greggio, L., 1986, *A&A*, 154, 279
 Matteucci, F., & Brocato, E., 1990, *ApJ*, 365, 539
 Matteucci, F., Romano, D., & Molaro, P., 1999, *A&A*, 341, 458
 McWilliam, A., & Rich, R.M., 1994, *ApJS*, 91, 749
 Mihalas, D., & Binney, J., 1981, *Galactic Astronomy* (second edition), (New York: W.H Freeman and Company)
 Minniti, D., 1996, *ApJ*, 459, 579
 Nakasato, N., Mori, M., & Nomoto, K., 1997, *ApJ*, 484, 608
 Nakasato, N., 2000, Ph.D. thesis, Department of Astronomy University of Tokyo
 Nakasato, N., Mori, M., & Nomoto, K., 2000, *ApJ*, 535, 776
 Navarro, J.F., & Steinmetz, M., 2000, *ApJ*, 538, 477
 Navarro, J.F., & White, S.D.M., 1993, *MNRAS*, 265, 271
 Navarro, J.F., & White, S.D.M., 1994, *MNRAS*, 267, 401
 Navarro, J.F., & Steinmetz, M., 2000, *ApJ*, 528, 607
 Nomoto, K., Yamaoka, H., Shigeyama, T., Kumagai, S., & Tsujimoto, T., 1994, in *Supernovae, Les Houces Session LIV*, eds. Bludman et al. (Amsterdam: North-Holland), 199
 Nomoto, K., Hashimoto, M., Tsujimoto, T., Thielemann, F.-K., Kishimoto, N., Kubo, Y., & Nakasato, N., 1997, *Nucl. Phys. A.*, 616, 79c
 Norman, C.A., Sellwood, J.A., & Hasan, H., 1996, *ApJ*, 462, 114
 Padmanabhan, T., *Structure formation in the universe* (Cambridge: Cambridge University Press)
 Patsis, P.A., & Athanassoula, E., 2000, *A&A*, 358, 45
 Raiteri, C.M., Villata, M., & Navarro, J.F., 1996, *A&A*, 315, 105
 Rich, R.M., 1988, *AJ*, 95, 828
 Rich, R.M., & McWilliam, A., 2000, in *Discoveries and Research Prospects from 8–10-Meter-Class Telescopes*, ed. J. Bergeron, SPIE vol. 4005, 150 (astro-ph/0005113)
 Samland, M., Hensler, G., & Theis, Ch., 1997, *ApJ*, 476, 544
 Sommer-Larsen, J., & Dolgov, A., 2001, *ApJ*, 608
 Springel, V., & Hernquist, L., 2002, *MNRAS*, 333, 649
 Steinmetz, M., 1996, *MNRAS*, 278, 1005
 Steinmetz, M., & Müller, E., 1994, *A&A*, 281, 97
 Steinmetz, M., & Müller, E., 1995, *MNRAS*, 276, 549
 Sugimoto, D., Chikada, Y., Makino, J., et al., 1990, *Nature*, 345, 33
 Sutherland, R.S., & Dopita, M.A., 1993, *ApJS*, 88, 253
 Thacker, R.J., & Couchman, H.M.P., 2000, *ApJ*, 545, 728
 Tsujimoto, T., Nomoto, K., Yoshii, Y., Hashimoto, M., Yanagida, S., & Thielemann, F.-K., 1996, *MNRAS*, 277, 945
 Umemura, M., Fukushige, T., Makino, J., Ebisuzaki, T., Sugimoto, D., Turner, E., & Loeb, A., 1993, *PASJ* 45, 311
 Yoshikawa, K., Jing, Y.P., & Suto, Y., 2000, *ApJ*, 535, 593
 Whitford, A.E., 1978, *ApJ*, 226, 777
 Wyse, R.F.G., 1999, *Cambridge Contemporary Astrophysics Series, The Formation of Galactic Bulges*, eds. Carolo, C.M., Ferguson, H.C., & Wyse, R.F.G. (Cambridge: Cambridge University Press), 195

TABLE 1
THE HEAVY ELEMENT YIELDS USED IN OUR CODE

	M ^a	Z	Fe	O
Type II SNe	14.0 $M_{\odot}$	2.54 $M_{\odot}$	0.091 $M_{\odot}$	1.80 $M_{\odot}$
Type Ia SNe	1.4 $M_{\odot}$	1.4 $M_{\odot}$	0.74 $M_{\odot}$	0.14 $M_{\odot}$

Note. — These data are taken from Tsujimoto et al. (1996) and Nomoto et al. (1997).
^aThe total ejected mass per one supernova explosion.

TABLE 2
KINEMATIC PROPERTIES OF THE MODEL GALAXIES.

C^a	$(\sigma_z/\sigma_x)^b$	spin (λ)	stellar mass ^c
0.1	0.59	0.70	$4.2 \times 10^{10} M_{\odot}$
0.5	0.71	0.52	$5.4 \times 10^{10} M_{\odot}$
1.0	0.77	0.44	$4.6 \times 10^{10} M_{\odot}$

Note. — ^a The star formation parameter.
^b The ratio between the X and Z axis velocity dispersion.
^c The stellar mass at $t = 5$ Gyr.




FIG. 1.— The star formation probability (P) as a function of density of gas. The solid and dashed lines correspond to $C = 1.0$ and $C = 0.1$, respectively.




FIG. 2.— The star formation rate (SFR) as a function of time for galaxies with $C = 0.1$ (the dotted line) and $C = 1.0$ (the solid line).




FIG. 3.— The Type Ia supernova rate (R_{SNIa}) as a function of time.




FIG. 4.— The schematic view of the feedback processes in our SPH code. “ $t(m)$ ” is the stellar lifetime as a function of main-sequence mass ($m/M_{\odot}$).




FIG. 5.— The projected STAR positions at $t = 5$ Gyr ($z \sim 1$) for the three components. From the left panel, the metal-poor STARS, the metal-rich and old STARS, and the young STARS are shown (see text). The size of the panels is $20\text{kpc} \times 20\text{kpc}$.




FIG. 6.— The star formation history for the bulge stars (solid line) and all stars (dotted line). The SFR divided by the total stellar mass of each population is shown as a function of time.




FIG. 7.— The face-on surface density of STARS at $t = 5$ Gyr. The crosses represent our model and the solid line is the fit to our model (see text).




FIG. 8.— The projected STAR position for the first 0.5 Gyr of the evolution. During $t = 0.4 - 0.5$ Gyr, two clumps merge together. The size of the panels is $10\text{kpc} \times 10\text{kpc}$.




FIG. 9.— The mean velocity and the velocity dispersion for the bulge stars as a function of the distance from the galactic center. The upper and lower lines are the average velocity and the velocity dispersion, respectively.




FIG. 10.— The x-direction velocity dispersion for the bulge stars as a function $[\text{Fe}/\text{H}]$. The dotted circles are the observational data in Minniti (1996) (the error bound in the observation is $\sim 8 - 14 \text{ km s}^{-1}$). Our model STARS are shown by the filled squares ($t_{\text{form}} < 1 \text{ Gyr}$) and the filled triangles ($t_{\text{form}} < 5 \text{ Gyr}$).




FIG. 11.— The calculated metallicity ($[\text{Fe}/\text{H}]$) distribution function of the bulge stars (solid line). The dashed line shows the observed metallicity distribution function of K-giant stars (McWilliam & Rich 1994). The distributions are normalized by its maximum value for comparison.




FIG. 12.— The calculated metallicity ($[\text{Fe}/\text{H}]$) distribution function of the bulge stars (solid line). The distributions are not normalized. The dashed and dotted lines correspond to the very old STARS with $t_{\text{form}} < 0.5$ Gyr and young STARS with $t_{\text{form}} \geq 0.5$ Gyr, respectively.




FIG. 13.— The calculated metallicity ($[O/Fe]$) distribution function of the bulge stars.




FIG. 14.— $[O/Fe]$ vs $[Fe/H]$ for ~ 800 STARS, which are randomly selected from the bulge stars in our model (filled circles). The dotted circles show preliminary observational data (Rich & McWilliam 2000).